

# Sexual violence towards females in Matara

Rathnaweera RHAJ

Department of Forensic Medicine, Faculty of Medicine, Karapitiya, Galle

\*Corresponding author: Tel: 0094-772969060. E-mail address: [ajithrathnaweera@yahoo.com](mailto:ajithrathnaweera@yahoo.com)

MLJSL. Vol 1. No 3. december. pp 08-13

---

## Abstract

### Introduction

Information on sexual violence towards females is inadequate. Some statistics are available, but it is uncertain how accurate they are, as many cases go unreported. Therefore an in depth study should be conducted to explore the real situation.

### Objective

Routine data pertaining to victims of alleged sexual violence were analyzed to document epidemiological and socio-economic data and to describe the medico-legal aspects of injuries.

### Materials and Methods

All the sexual violence cases referred to the JMO's office, General hospital, Matara from 1<sup>st</sup> of January 2012 to 1<sup>st</sup> of December 2012 were retrospectively analyzed.

### Results

During the study period, 260 victims were referred for medico-legal examination. The age of the victims ranged from 4 years to 62 years. 177 (68%) victims were between 11 to 16 years of age. Majority (90%) of the victims belonged to lower socio-economic classes. 210 (80%) had not studied beyond grade 10. The alleged incident had taken place at the victims' house in 110 (42%) cases. In 148 (57%) cases, the victim had given "consent" for the alleged act. 22 (8%) were pregnant at the time of examination. Out of the population 100(38%) had hymenal tears but out of that, only 11 had fresh tears. Extra genital injuries were present only in 4 (1.5%) cases. The assailant was the boy friend in 149 (57%) cases. Seventy (27%) cases were reported from Matara police area followed by Akuressa 43 (17%).

### Conclusions

Majority were children below 16 years of age belonging to lower socio-economic group. As they were below 16 years, this amounted to statutory rape. Majority had given "consent" for the alleged act and this could be a reason for lack of extragenital injuries together with delayed presentation. This data highlights the magnitude of juvenile sexual abuse especially by the older boy friends and other known people.

**Keywords:** *Sexual violence, hymenal tears, known assailants, statutory rape*

---

## Introduction

Sexual violence refers to sexual activity where consent is not obtained or freely given [1]. Coercive strategies used in sexual violence may be physical, verbal, or psychological. Sexual violence is a serious public health problem that threatens the health and well-being of individuals across the world. Victims of sexual violence may experience an increased risk of acute and chronic health problems, including injuries resulting from the sexual assault and subsequent health problems and health risk behaviours. For perpetrators, arrest and conviction may result in social and economic costs related to incarceration or community notification and registration laws. The economic burden of sexual violence victimization including medical and mental health care, loss of work, and impact on quality of life was estimated in 1996 to be approximately \$126 billion annually in the United States [1].

Information on sexual violence towards females is inadequate. Some statistics are available, but it is uncertain how accurate they are, as many cases go unreported. The National Intimate Partner and Sexual Violence Survey found that 1 in 5 women and 1 in 71 men experience sexual violence during their lifetimes [2]. In Swedish population-based surveys, 12% to 34% of Swedish women have been exposed to some kind of sexual assault during their lifetime [3]. A study conducted in the early 1990s among 899 pre-university and undergraduate students in Sri Lanka revealed that 18% of boys and 4.5% of girls had been sexually abused in childhood [4]. Most of the perpetrators were known parties such as relatives, neighbors, teachers and priests. As indicated by the above study, an in depth study should be conducted to explore the real situation.

## Materials and Methods

All the sexual violence cases referred to the JMO's office, General hospital, Matara from 1<sup>st</sup>

of January 2012 to 1<sup>st</sup> of December 2012 were retrospectively analyzed. The medico legal records of 301 consecutive sexually assaulted cases were evaluated. Of the 301 cases, 21 cases were excluded due to incomplete documentation. Sexual assault cases against males (20) were also excluded. Routine data pertaining to female victims of alleged sexual violence were analyzed to document epidemiological and socio-economic data and to describe the medico-legal aspects of injuries. The different types of assault were defined as penetration (with penis, hand, or foreign object) vaginal, oral, anal or a combination of the three, alternatively others when no penetration was involved but other kinds of touching or kissing had occurred (involving genitals or body parts). Violence was defined as hitting, kicking, holding, attempts of strangulation, or by other means.

Genital injuries were defined when swelling (oedematous tissues), redness (erythematous skin), ecchymoses (skin or mucous membrane discolorations, known as bruising), tears (breaks in tissue including fissures, cracks, cuts and lacerations), and several (more than one of the above mentioned injuries) were observed. The different genital injury locations are defined as vulva, introitus, labia minora, posterior fourchette, hymen, and vagina.

## Results

During the study period, 277 female victims were referred for medico-legal examination with an alleged history of sexual abuse. Out of that 17 cases were excluded as the victims did not recall or were unwilling to relate assault details. The age of the victims ranged from 4 years to 62 years (Table 01). 177 (68%) victims were between 11 to 16 years of age.

Majority (90%) of the victims belonged to lower socio-economic classes whereas 8% to middle classes and only 2% to higher classes.

210 (80%) had not studied beyond grade 10.

The alleged incident had taken place at the victims' house in 110 (42%) cases (Table 02). In

148 (57%) cases, the victim had given the “consent” for the alleged act. 22 (8%) were pregnant at the time of examination.

**Table:1** Age distribution of the victims

Age (Years)	No
1-5	08 (3%)
6-10	17 (6.5%)
11-16	177 (68%)
17-20	30 (11.5%)
21-25	09 (3.4%)
26-30	04 (1.5%)
31-35	03 (1.1%)
36-40	06 (2.3%)
41-45	02 (0.7%)
46-50	03 (1.1%)
51-55	00
56-60	00
61-65	01 (0.3%)

**Table 02 :** Alleged place where incident had taken place

Place	No
Victim’s home	110 (42%)
Assailant’s home	35 (13%)
Hotel/Guest house	34 (13%)
Friend’s home	26 (10%)
Relative’s home	26 (10%)
Neighbour’s home	28 (3%)
Deserted place	15 (5.7%)
Temple	02 (0.7%)
School	02 (0.7%)
Film hall	02 (0.7%)

The alleged abuse was vaginal intercourse in 225 cases followed by kissing in 153 and intercrural intercourse in 91 (Table 03).

Annular hymen 125 (48%) was the commonest type of hymen found among the population followed by fimbriated 67(25%) and crescentic 47(18%). Five (1.9%) had attenuated hymen and 16 (6%) had carunculae multiformes.

Ano-genital injuries were present in 154 victims (Table 04).

**Table 03:** Allegation

Type of abuse	No
Kissing	153
Fondling	36
Intercrural intercourse	91
Oral intercourse	45
Vaginal intercourse	225
Anal intercourse	11
Combination	241

**Table 04:** Medical evidence

Type of injury	No
Hymenal tears	100
Reddening of hymen	25
Fourchette tear	31
Dilated orifice	71
Loss of vaginal rugae	61
Anal injuries	01
Reddening of Labia	16
Combination	131

Out of the population 100(38%) had hymenal tears and out of that, only 11 had fresh tears (Table 05).

**Table 05:** Nature of the injury

Type of tear	No
Fresh	11
Partially healed	14
Old healed	75

6 o’ clock position was the commonest place for hymenal tear followed by 9 o’ clock and 3’o clock.

Hymenal tears were observed in 53 annular hymens, 34 fimbriated hymens and 13 crescentic hymens. For annular hymens, 6’ o clock was the commonest place of tear, for fimbriated hymens, 3’ o clock and for crescentic hymens, 6’ o clock was the commonest place of tear (Table 06).

Extragenital injuries were present only in 4 (1.5%) cases and in all 4 cases the assailant was an unknown person.

Abrasion was the Commonest type of injury found. Face was the commonest place to be injured followed by upper limbs and lower

limbs.

**Table 6:** location of tear in different types of hymen

Location	Annular hymen	Fimbriated hymen	Crescentic hymen
3 o' clock	13	31	03
5 o' clock	08	11	03
6 o' clock	48	18	13
7 o' clock	06	03	06
9 o' clock	15	29	04
11 o' clock	05	06	04

The assailant was the boy friend in 149 (57%) cases and only in 8 (%), the assailant was an unknown person (Table 7).

**Table 7:** Assailant

Assailant	No
Boy friend	149 (57%)
Father	10 (3.8%)
Relative	12 (4.6%)
Priest	02 (0.7%)
School teacher	02 (0.7%)
Known person	77 (30%)
Unknown person	08 (3.0%)

The assailants were aged from 14 years to 57 years (Table 8). Majority of the assailants were aged between 21-25 years 103(40%).

**Table 8:** Age of the assailant

Age of the assailant	No
>15	02 (0.7%)
16-20	89 (33%)
21-25	103 (40%)
26-30	24 (9%)
31-35	06 (2.3%)
36-40	15 (5.7%)
41-45	06 (2.3%)
46-50	10 (3.8%)
51-55	04 (1.5%)
56-60	03 (1.1%)

Seventy (27%) cases were reported from Matara police area followed by Akuressa 43 (17%).

## Discussion

Several studies have been published during the recent years examining genital and nongenital injuries in female sexual assault victims. Sexual assaults by strangers have been reported as more violent and associated with more injuries than assaults by a person known to the victim [5 - 9]. This is in keeping with findings of this study as all the victims who had extragenital injuries had been abused by an unknown person.

Majority of the victims were children below 16 years of age belonging to lower socio-economic group. As they were below 16 years, this had amounted to statutory rape. Extragenital injuries were present only in 4 (1.5%) cases. Majority had given "consent" for the alleged act and this could be a reason for lack of extragenital injuries.

Seventy (27%) cases were reported from Matara police area followed by Akuressa 43 (17%). The reason for these high percentages should be studied separately. The alleged incident took place at the victim's house in majority of cases. Only in very few cases it took place at a deserted place and the assailant was an unknown person. This data highlights the magnitude of juvenile sexual abuse especially by the older boy friends and other known people.

The victim's relationship to the assailant is consistent with earlier international studies [10,11] and the majority of all assaults were committed by a person known to the victim.

Sexual assaults committed by an unknown person comprise 03% in this study population, which is somewhat lower when compared with studies from the United Kingdom and the United States [5,12].

The alleged abuse was most frequently vaginal intercourse followed by kissing and intercrural intercourse. Ano-genital injuries were present

in 154 victims. Out of those who complained of vaginal intercourse, 100 had hymenal tears and only 11 had fresh tears. For annular hymens, 6 o'clock was the commonest place of tear, for fimbriated hymens, 3 o'clock and 9 o'clock were the commonest place of tear and for crescentic hymens, 6 o'clock was the commonest place of tear. This type of pattern has not been described in the literature and further studies should be carried out to get further information in this regard.

Our results, establishing that only a minority of female sexual assault victims have clinical evidence of fresh genital injuries, are supported by previous studies [6, 10, and 12].

The genitoanal area has a remarkable healing capacity resulting in injuries disappearing quickly. One of the most important factors in the forensic examination is therefore the time lapse between the sexual assault and the examination. An examination after 72 hours will not show minor lacerations that might have been present earlier.

This study has some limitations and strengths that need to be mentioned. The study is retrospective, and the choice of variables was limited. Data collection was dependent on victim's report, and it is not clear how reliable it is to collect data within the context of a forensic examination shortly after a sexual assault because of peritraumatic dissociation. Some women did not recall or were unwilling to relate assault details. Although no more than 17 of responses were missing for any one variable, it is possible that these missing data could have affected study findings. Data are collected from one hospital that in most cases probably would reduce generalizability, but as all sexual assault victim care in the Matara district are centralized to one unit, these are the circumstances that give strength to the study.

## Conclusions

Some of the police areas have a higher percentage of cases reported than the others and the reason for these high percentages

should be studied separately. Majority of the victims were children below 16 years of age belonging to lower socio-economic group. Majority had given "consent" for the alleged act and this could be a reason for lack of extragenital injuries together with delayed presentation. This data highlights the magnitude of juvenile sexual abuse especially by the older boy friends and other known people. Sexual assaults committed by an unknown person comprise a low percentage in this study when compared with studies from the United Kingdom and the United States. A different pattern of hymenal injuries were observed for different types of hymens and this type of pattern has not been described in the literature and further studies should be carried out to get further information in this regard.

## References

1. Tharp AT, DeGue S, Valle LA, Brookmeyer KA, Massetti GM. A Systematic Qualitative Review of Risk and Protective Factors for Sexual Violence Perpetration. Trauma, violence and abuse. Published online on 27 December 2012. Available at <http://tva.sagepub.com/content/early/2012/12/27/1524838012470031>
2. Black MC, Basile KC, Breiding MJ, Smith SG, Walters ML, Merrick MT, Chen J, & Stevens MR (2011). The National Intimate Partner and Sexual Violence Survey (NISVS): 2010 Summary Report. Atlanta, GA: Centers for Disease Control and Prevention.
3. Möller A.S, Bäckström T, Söndergaard HP, Helström L. Patterns of Injury and Reported Violence Depending on Relationship to Assailant in Female Swedish Sexual Assault Victims. *Journal of Interpersonal Violence* 2012;27(16): 3131–3148
4. De Silva H. Second country report on the implementation of the convention on the rights of the child. Department of Probation and Childcare Services, Ministry of Social Services Sri Lanka 1998.

5. Jones, JS, Wynn, BN, Kroeze B, Dunnuck C, Rossman L. Comparison of sexual assaults by strangers versus known assailants in a community-based population. *American Journal of Emergency Medicine* 2004; 22, 454-459.
6. Palmer CM, McNulty AM, Déste C, Donovan B. Genital injuries in women reporting sexual assault. *Sex Health* 2004; 1, 55-59.
7. Slaughter L, Brown CR. Patterns of genital injury in female sexual assault victims. *American Journal of Obstetrics & Gynecology* 1997; 176, 609-616.
8. Stermac LE, Du Mont JA, Kalemba V. Comparison of sexual assaults by strangers and known assailants in an urban population of women. *Canadian Medical Association Journal* 1995; 153, 1089-1094.
9. Sugar NF, Fine DN, Eckert LO. Physical injury after sexual assault: Findings of large case series. *American Journal of Obstetrics & Gynecology* 2004; 190, 71-76.
10. Grossin C, Sibille I, Lorin de la Grandmaison, G, Banasr A, Brion F, & Durigon M. Analysis of 418 cases of sexual assault. *Forensic Science International* 2003; 131, 125-130.
11. Logan TK, Cole J, Capillo A. Differential characteristics of intimate partner: Acquaintance, and stranger rape survivors examined by a sexual assault nurse examiner. *Journal of Interpersonal Violence* 2007; 22, 1066-1076.
12. Maguire W, Goodall E, Moore T. Injury in adult female sexual assault complainants and related factors. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2009; 42, 149-153.